

E-palene Cream
(A d a p a l e n e)

15g
0.1%

ای-پالین
(ایڈاپالین)

COMPOSITION

E-palene 0.1% cream: Each gm cream contains Adapalene 1 mg.

PHARMACOLOGY

Adapalene acts on retinoid receptors that are commonly found in the skin of face, back and chest. Biochemical and pharmacological studies have demonstrated that Adapalene is a modulator of cellular differentiation, keratinization, and inflammatory processes, all of that represent important features in the pathology of acne vulgaris. Adapalene binds with specific retinoic acid nuclear receptors that normalize the differentiation of follicular epithelial cells resulting in decreased microcomedone formation. Absorption of Adapalene through human skin is low.

INDICATION

E-palene 0.1% cream a gel is indicated for topical treatment of acne vulgaris.

DOSAGE AND ADMINISTRATION

- **E-palene** 0.1% cream: E-palene 0.1% cream should be applied to the affected areas of skin, once daily at night-time
- A thin film of cream should be applied to the skin areas where lesions present, using enough to cover the entire affected areas lightly
- For detail information, please see the enclosed leaflet

CONTRAINDICATION

Adapalene should not be administered to individuals who are hypersensitive to Adapalene or any of its components.

PRECAUTION

Adapalene should not be applied to cuts, abrasions, eczematous or sunburned skin.

SIDE EFFECT

Erythema, scaling, dryness, pruritus, burning sensation, skin irritation, stinging unburn, acne flares, etc. are commonly seen during the first month of therapy but usually lessen with continued use of the medication.

DRUG INTERACTION

- Concomitant use of other potentially irritating topical products (medicated or abrasive soaps and cleansers, soaps and cosmetics that have a strong drying effect, products with high concentrations of alcohol, astringents, spices or lime) should be approached with caution
- Exercise particular caution in using preparations containing sulfur, resorcinol or salicylic acid in combination with Adapalene
- If any of these preparations have been used, it is advisable not to start therapy with Adapalene until the effects of such preparations in skin have subsided. If combined use of both medications is important, it is better to use in two different times

USE IN PREGNANCY AND LACTATION

Use Adapalene during pregnancy only if the potential benefit justifies the potential risk to the fetus. It is not known whether this drug is excreted in breast milk.

Exercise caution when administering Adapalene to a nursing mother.

USE IN CHILDREN

Safety and effectiveness in children below 12 years of age have not been established.

STORAGE

Store in a cool and dry place (20°C to 25°C). Do not freeze.

HOW SUPPLIED

E-palene 0.1% cream: Each pack has a Aluminum tube containing 15 gm cream.

INSTRUCTIONS:

Use as directed by the physician. Keep out of the reach of children.
Protect from light & freezing. Store at temperature between 20°C to 25°C.

ہدایات: ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔
بچوں کی پہنچ سے دور رکھیں۔ روشنی اور ٹھنڈ ہونے سے بچائیں۔
۲۰ سے ۲۵ ڈگری سینٹی گریڈ درجہ حرارت کے درمیان رکھیں۔

Manufactured by:
ISO Certified Company
Medera Pharmaceuticals (PVT) LTD.
Plot No 2 street No N-4, National Industrial Zone
Rawat-Pakistan